Efficient reduction of CO₂ using a novel carbonic anhydrase producing *Corynebacterium flavescens*

Tanvi Sharma, Ashok Kumar[†]

Department of Biotechnology and Bioinformatics, Jaypee University of Information Technology, Waknaghat, Solan, India

ABSTRACT

Emission of greenhouse gases into the atmosphere by human activities leads to global warming. To reduce the level of CO₂ the bio-catalytic properties of microbial carbonic anhydrase (CA) can be exploited. The present study aimed to isolate CA producing bacteria from cow saliva. After thorough screening for CA activity in the bacterial cultures, ten isolates were selected. Out of ten bacterial isolates, T5 isolate showed the highest CA activity (83.92 U/mL) and the isolate was identified as *Corynebacterium flavescens* using 16s rRNA analysis. Various production parameters for the optimum production of CA were optimized. During the optimization, incubation temperature 40°C, agitation speed 120 rpm, and inoculum volume 4%v/v was found to be optimum for CA production. The optimum reaction pH, reaction time and temperature were 7, 10 min and 35°C, respectively. The crude enzyme was tested for the conversion of CO₂ into the calcium carbonate (CaCO₃) under controlled conditions. The CO₂ conversion efficacy of crude CA was observed to be ~45 mg CaCO₃/mg protein. The synthesized CaCO₃ was analyzed using scanning electron microscopy and X-ray diffraction techniques for particle size, morphology and elemental structure. Calcite precipitation by bacterial CA makes it a potential candidate to be effectively employed in biomimetic CO₂ sequestration.

Keywords: Carbonic anhydrase, CO₂ sequestration, *Corynebacterium flavescens*, SEM

1. Introduction

Global warming is one of the serious issues due to excessive emission of carbon dioxide (CO₂) into the atmosphere. To reduce CO₂ emission, numerous government and non-governmental organizations endorse to switch from fossil fuel power to clean energy sources to protect the atmosphere from the terrible effects of global warming [1-3]. Furthermore, worldwide research is going on for finding a potent and eco-friendly method that directly converts industrial emitted CO₂ into a useful product. Some of the algae, bacteria, and cyanobacteria play a vital role in alleviating the increasing level of CO₂ using their carboxylating enzyme [4, 5]. Recently, these carboxylating enzymes are getting attention due to its role in CO₂ sequestration [1, 6]. However, algae have good potential for CO₂ fixation, but it has a few disadvantages such as light requirement [7]. Hence, it is advantageous to isolate a CO₂ fixing bacteria or enzyme which does not require continuous light for its growth and activity [8].

CA is one of the fastest known biocatalysts and involved in the CO₂ sequestration. CA contains zinc in its active center and converts CO₂ to bicarbonates, which can be further converted to

CaCO₃ in the presence of calcium ion. The application of CA in the conversion of CO₂ from flue gas into thermodynamically stable, environmentally safe calcium carbonate offers several advantages [9]. First of all, the process is extremely fast and take place near ambient condition. CaCO₃ formed during CO₂ conversion can be utilized in the preparation of white pigment, cement, antacids and others [10]. Thus, a new biomimetic approach using CA has been found to be viable for fixing a huge quantity of CO₂ into CaCO₃. CA catalyzes several other hydrolytic reactions too, including hydration of urea, carboxylic acid, halides, and hydrolyzable substrates [11, 12]. The CA catalyze the hydration of CO₂ in two steps ping-pong mechanism. Its active site has divalent metal ions generally zinc in a tetrahedral confirmation, containing three amino acid as ligands and hydroxide ions coordinating the metal [13]. CA is present in eukaryotes and prokaryotes, where it takes part in various physiological functions such as acid-base balance, hemostasis, respiration, and photosynthetic CO₂ fixation [14]. Although CA is present ubiquitously but microbial CA has received more attention due to ease of their production and applications. Till now, CA has been classified into five (α , β , γ , δ and ϵ) different

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2021 Korean Society of Environmental Engineers

Received April 21, 2020 Accepted July 08, 2020

[†] Corresponding author

Email: ashok.nadda09@gmail.com

Tel: +01792-239351-353 Fax: +01792-239351

ORCID: 0000-0001-9192-0774

Fig. 1. Schematic representation of isolation, optimization and CaCO_3 production using carbonic anhydrase.

classes [15]. The first studied, class is the α -CA as it plays a crucial role in human pathology and drug targeting. However, the other classes take part in the conversion of gaseous CO_2 [16]. The β class exists as a dimer, tetramer and hexamer and mostly present in algae, bacteria, and archaea. The γ class is mostly found in archaea and exists as a trimer [17]. This enzyme is either present extracellularly or inside (intracellular) the cytoplasm in the bacteria [18].

Recent studies suggest that CA is widespread in bacteria and archaea domain, indicates that this enzyme has a more extensive and vital role in prokaryotic physiology [19]. CA producing bacteria were isolated from various sources such as deep sea-water, alkaline soil, mangroves soil, and seashore water [20]. Moreover, the optimization of various parameters is essential to enhance the total yield, maximal activity of the enzyme and to reduce the process cost. The high enzyme titer has been attained by changing the ratio of various media ingredients that are influencing the growth and enzyme production in bacteria. The optimization is greatly influenced by various physicochemical and nutritional factors such as temperature, agitation rate, pH, media components, inoculum volume, *etc.* Thus, optimizing the best conditions for maximum enzyme production is still vital and essential in biocatalytic transformations.

In the present study, CA producing bacteria was isolated from cow saliva and the parameters were optimized for the production of CA in bulk amount. The application of crude CA has been studied for the conversion of CO_2 into calcium carbonates (Fig. 1). As per our knowledge, this is the first report on the isolation of CA producing bacteria from cow saliva. In the current global warming scenario, this work might be helpful in CO_2 conversion and environmental amelioration. Here, we demonstrate that isolated bacteria are the green biocatalyst for the conversion of CO_2 into calcium carbonates.

2. Methods

2.1. Isolation of a Bacterial Strain

CA is a ubiquitous enzyme present in all plants, mammalian tissues, algae, and bacteria [19]. In the present study, the sample was collected from six years old cow saliva from District Mandi, Himachal Pradesh, India with the help of sterilized bud (Fig. 2). The saliva was serially diluted and then 100 μL of culture was transferred

Fig. 2. Flowchart summarising the overall research methodology.

onto nutrient agar having 3mM *p*-nitrophenyl acetate (*p*-NPA), incubated at 30°C for 48 h. The appearance of the intense yellow colour colonies indicated the production of carbonic anhydrase [21]. The bacterial isolates, which utilized *p*-NPA on agar plates, were further screened for CA activity in nutrient broth. For CA activity assay, the seed culture was prepared aseptically in 50 mL nutrient broth containing a loopful culture of T5 isolate and incubated at 30°C for 24 h. For the production of CA, the nutrient broth was used as production media and the seed culture 2% v/v was used as inoculum for 50 mL production medium. The inoculated production medium was incubated at 30°C for 36 h at 120 rpm. The broth was centrifuged at 10,000 rpm for 10 min after 36 h of incubation. Then pellet was suspended in phosphate buffer (pH 7.0) and homogenized well. The suspension was sonicated using 20 kHz frequency for 5 min at 4°C and centrifuged at 10,000 rpm for 20 min. The supernatant was used as crude enzyme extract.

2.2. Enzyme Assay

CA activity was assayed in the culture broth using a previously reported method by measuring the micromole of *p*-nitrophenol released from *p*-NPA [22]. To 825 μ L of Phosphate buffer (50 mM, pH 7.5), 175 μ L of the substrate stock solution (*p*-NPA, 10 mM in isopropanol) was added. Then, the reaction mixture was incubated in a water bath at 37°C for 5 min. Subsequently to initiate the reaction 25 μ L of an enzyme was added. The reaction mixture was again incubated at 37°C in a water bath for 5 min [23]. The amount of *p*-nitrophenol released was measured at 410 nm after 5-min incubation using a microplate reader (Thermo-scientific, Multiskan™ FC Microplate Photometer) [24]. Based on CA activity, a potent isolate T5 was selected for all subsequent studies. The colonies of T5 isolate was subjected to gram's staining and observed at 100 X magnification under a compound microscope. The biochemical characterization of T5 isolate was conducted according to Bergey's Manual of Systemic Bacteriology [25].

2.3. Identification of Bacterial Strain

The bacterial strain was identified by 16S rRNA sequencing by Bioreserve Biotechnologies Pvt. Ltd., Hyderabad, India [26]. The nucleotide sequences obtained from 16s rRNA sequencing were subjected to a homology search using BLAST and aligned using a MEGA X software. Then, the phylogenetic tree was made using the neighbour-joining method.

2.4. Optimization of Production Parameter for CA Producing Bacteria

Luria broth, nutrient broth, peptone broth, minimal salt media, muller Hinton broth, and basal salt media were used for enzyme production. Luria broth, nutrient broth, and muller hinton broth were procured from Hi Media, Mumbai. The composition of other media used for enzyme production was peptone broth (g/L): beef extract 3.0, glucose 1.0, NaCl 5.0, peptone 5.0, CaCO₃ 6.0; minimal salt media (g/L): KH₂PO₄ 6.8, MgSO₄ 0.2, Na₂HPO₄ 7.8, ZnCl₂ 0.02, ZnSO₄·7H₂O 0.05, NaNO₃ 0.085; and basal salt media (g/L): sucrose 5.0, MgSO₄·7H₂O 0.5, Na₂HPO₄ 2.0, FeCl₃·6H₂O 0.005, CaCO₃ 0.1. The effect of temperature on the CA production was studied by

incubating production culture at various temperatures ranging from 20 to 60°C. To determine the effect of inoculum volume (%v/v) on CA production, production media was inoculated with a (2-12 %v/v) culture of T5 isolate and incubated at 40°C. The optimum agitation rate for CA production was determined by incubating the production media at 40°C at varying agitation speeds (80, 100, 120, 140, 160, and 180 rpm). The enzyme activity was determined for each incubated production media sample using standard activity assay [23].

2.5. Optimization of Reaction Parameter for CA Producing Bacteria

For optimizing reaction conditions of CA effect of reaction time (2-18 min), different buffer systems (citrate buffer, potassium phosphate buffer, and Tris-HCl buffer, 50 mM), reaction temperature (25-60°C), different metal ions (Na⁺, K⁺, Ca²⁺, Zn²⁺, Mg²⁺, Fe³⁺, and Al³⁺) and organic solvent (ethanol, isopropanol, ethanediol, *n*-butanol, propandiol, and, acetonitrile) was studied on CA activity.

2.6. Enzymatic Conversion of Carbon Dioxide into Calcite

The potential of CA for CO₂ conversion was determined as described by Giri et al. [27], with minor modifications. The precipitation of CO₂ into calcium carbonate (CaCO₃) was carried out in a 50 mL total reaction mixture containing 2 mL Tris buffer (1M, pH 8), 23 mL of 2 (%v/v) calcium chloride solution, 23 mL of CO₂ saturated water and 2 mL (0.5 mg/mL) of an enzyme in phosphate buffer (50 mM, pH 7). The reaction was performed for 10 min at room temperature. Bovine serum albumin (BSA) was used as a control. The precipitate formed after 10 min was recovered by centrifugation. The sediment of precipitates was lyophilized overnight to obtain the dry powder. Then, CaCO₃ precipitates were weighed to determine the amount of carbonate deposited during the enzymatic reaction.

2.7. Instrumental Analysis of CaCO₃

The CaCO₃ precipitates were lyophilized to get dry powder of CaCO₃. The CaCO₃ precipitates were scanned in the range of 400-4,000 cm⁻¹. Compositions of the precipitated solid crystals were examined by using X-ray diffraction (XRD). To determine crystal morphology Scanning Electron Microscopy (SEM) was performed by using JSM7401F (JEOL).

3. Results and Discussion

3.1. Isolation of CA Producing Bacteria

The ten bacterial isolates from a cow saliva sample, which efficiently utilized *p*-NPA on agar plates, were selected for secondary screening and cultivated in nutrient broth to detect CA activity. *p*-NPA plates assay is specific and based on the formation of yellow-colored (*p*-NP) product formed around the colonies. Among these bacterial isolate T5, showed maximum CA activity, and selected for further work (Fig. 3 (a)). The T5 isolate was rod-shaped and gram-positive bacilli (Fig. 3 (b)). And the results of the biochemical analysis show Voges-Proskauer positive, nitrate reduction positive, glucose and sucrose positive. The presence of CA in mammalian saliva

Fig. 3. (a) Selection of bacterial isolate on the basis of CA activity. (b) Microscopic view of bacterial isolate T5.

Fig. 4. (a) Gel image of 16S rDNA amplicon [Lane 1: 16S rDNA amplicon and Lane 2: DNA markers]. (b) Phylogenetic tree of bacterial T5 isolate.

is reported several decades ago. Yoshimura et al. [28] suggested that the role of CA in mammalian saliva is to regulate saliva pH. Furthermore, the screening of microorganisms from cow saliva for CA production is the rapid and cost-effective method. Jaya et al. [29] reported an isolate *Bacillus safensis* from the water sample that produced CA. Similarly, Kupriyanova et al. [30] reported that the presence of CA in *Microcoleus cathonoplastes*, isolated from the soda lakes in Russia. Various pathogenic bacteria such as *Yersinia pseudotuberculosis* and *Listeria monocytogenes* isolated from soil and water were reported to produce CA [31]. Microorganisms need CO₂ and bicarbonate in their metabolic activities and CA plays a vital role to meet these demands.

3.2. Identification of Strain

The genomic DNA was isolated from bacterial isolate T5 and its

DNA analysis was done by agarose gel electrophoresis (Fig. 4 (a)). The 16s rRNA was amplified from genomic DNA using universal primers. Then bacterial strain was identified as *Corynebacterium flavescens* by 16S rRNA sequencing based on phylogenetic analysis and nucleotide homology. The phylogenetic tree was made by using a neighbour tree joining method showing maximum sequence homology of 99.42% and 99.27% with *Corynebacterium flavescens* HBUM07012 and *Corynebacterium flavescens* EHFS1S12Hc, respectively (Fig. 4 (b)). The gene accession number of the organism was MN982752 deposited in NCBI.

3.3. Optimization of Production Conditions for CA

Optimization of production and reaction parameters is widely known to rise the production significantly. In the present study, out of six media, the maximum CA activity (84.99 U/mL) was

Fig. 5. (a) Effect of different media (Luria broth [LB], nutrient broth [NB], Peptone broth [PB], minimal salt media [MS], basal salt media [BS], muller Hinton broth [MB]) on the production of carbonic anhydrase was determined, with activity at nutrient broth as 100%. (b) Effect of incubation temperature on carbonic anhydrase production was determined, with activity at 40°C as 100%. (c) Effect of inoculum volume (2, 4, 6, 8, 10, 12% v/v) on CA production was calculated, with activity at 4% v/v as 100%. (d) Effect of agitation rate on CA production was determined, with activity at 120 rpm as 100%.

observed in nutrient broth media (Fig. 5 (a)). In contrast, to the present findings, Sharma et al. [23] reported maximum CA production from *Pseudomonas fragi* in peptone broth. Also, in a previous study, the nutrient broth was reported as the best medium for the production of CA from *Nocardia opacensis* [20]. Thus, the nutrient broth media was selected for further studies. The temperature of incubation significantly affects the production of microbial enzymes and their activity. The maximum production of CA with residual activity 85.59 U/mL was observed at 40°C. Furthermore, a decrease in the CA activity was noticed with a further increase in temperature (Fig. 5 (b)) [24]. The decrease in activity might be attributed to the denaturation of protein at a higher temperature. Similarly, Sharma et al. [23], reported the optimum CA production was observed at 40°C. In a recent study, *Bacillus safensis* was observed to show optimum activity at 40°C [29]. In another study, CA from *Citrobacter freundii* was found to be active at 37°C [27]. However, the optimum temperature for CA production from *Methanobacterium thermoautotrophicum* was found to be 75°C [22]. These results showed that CA production has its own temperature optima and is species-specific which favours maximum CA production.

The volume of inoculum also affected the production and activity of enzymes from the isolated bacterial strain. An increase in CA

production was observed with an increase in an inoculum volume up to 4% v/v then it started decreasing (Fig. 5 (c)). Another study showed that 1.5% inoculum size was optimum for CA production from *Aeribacillus pallidus* [32]. Usually increased inoculum volume improves the growth of bacteria up to a certain level and after that bacterial growth starts decreasing due to nutritional limitation. Furthermore, lower inoculum volume has a low number of bacterial cells in medium and this requires a longer time to grow. This might lead to the accumulation of toxic compounds which ultimately decreased enzyme production [33]. However, some studies reveal that an increase in inoculum size results in more enzyme production and *vice versa*. The maximum CA activity was observed at an agitation speed of 120 rpm *i.e.*, 86.50 U/mL (Fig. 5 (d)). The CA activity was increased by increasing the agitation rate up to 120 rpm. But the high agitation rate also resulted in decreased activity due to mechanical disruption of proteins. In a previous study, carried out by Zhang et al. [34], showed that 150 rpm most appropriate agitation rate for the production of CA whereas CA from *Aeribacillus pallidus* gave maximum activity at 200 rpm [32]. The key role of agitation in the fermentation broth is to ensure adequate mixing of oxygen that eventually becomes available for the growth of bacteria in the dissolved form of oxygen. Also, microbial enzyme production

Fig. 6. (a) Effect of pH on activity. The enzyme activity at different pH (sodium citrate buffer of pH 4.0-5.5, Phosphate buffer of pH 6.0-7.5 and, Tris-HCl buffer of pH 8.0-9.5) was determined, with the activity at pH 7 as 100%. (b) The effect of different temperatures (25-60°C) on the activity was measured at pH 7.0, with the activity at 35°C as 100%. (c) The effect of reaction time on the activity was measured with the activity at 10 min as 100%. (d) Effect of organic solvents on CA activity was determined.

is observed to be good under continuous shaking conditions as compared to static conditions [35].

3.4. Optimization of Reaction Parameters for CA

The effect of pH on CA activity was determined and optimum pH for maximum CA activity was 7 in phosphate buffer (Fig. 6 (a)). A decrease in activity at high and low pH generally results in loss of enzyme activity due to disruption of three-dimensional structure and alteration in amino acid residues present in the active center of an enzyme. In another study, CA from *Bacillus sp.* gave maximum activity at pH 8 whereas CA from *Lactobacillus* showed optimum activity at pH 6 [36, 37]. However, CA from *Helicobacter pylori* showed optimum activity in an acidic environment and high acid tolerance [38]. The effect of reaction temperature on the CA activity was studied. The result presented in (Fig. 6 (b)) showed that the optimum temperature for CA activity was 35°C after that decline in CA activity was observed. In the previous study, the optimum temperature for purified CA was found to be 60°C [36]. Mostly, CA isolated from human erythrocyte and bovine shows CA activity in the temperature range of 35-40°C and in the pH range of 6.5-7.5 [39]. The optimum reaction time for CA activity was observed after 10 min of incubation (88.55 U/mL) (Fig. 6 (c)). After 10 min decrease in enzyme activity was observed which may be due to

the inhibition of product and denaturation of the enzyme. All these facts indicate the functional diversity of CA and the capability of this enzyme to perform diverse roles in the organisms living in extreme environments.

Table 1. Activity of Bacterial CA in The Presence of Selected Metal Ions

Metal ion	Concentration (mM)	Relative Activity (%)
NaCl	1	87.43
	5	82.16
KCl	1	66.76
	5	70.69
MgCl ₂	1	54.45
	5	48.88
ZnCl ₂	1	103.75
	5	106.00
CaCl ₂	1	100.42
	5	99.30
FeCl ₃	1	101.64
	5	103.71
AlCl ₃	1	44.41
	5	37.47

Fig. 7. SEM image of CaCO_3 precipitate (a) Spherical vaterite formed in control (b) Spherical vaterite and rhombohedral calcite crystal formed in presence of enzyme (c) XRD analyses of CaCO_3 precipitates.

Out of various organic solvent used, all of them decrease CA activity (Fig. 6 (d)). This can be credited to conformational changes in enzyme active sites that are responsible for the decrease in enzyme activity [40]. Furthermore, the effect of metal ion on CA

activity at concentrations 1mM and 5mM was studied (Table 1). The presence of metal ion in the reaction medium either inhibits or promotes the enzymatic reaction. CA activity was found to be increased in presence of Zn^{2+} and Fe^{3+} ion this might be due to reason these metal ions are present in the active site of CA, they could have a stabilization effect thus enhancing the CA activity. In contrast, CA activity was not affected by the presence of Ca^{2+} ion indicates that enzyme could perform in the presence of calcium ion. Whereas, CA activity was inhibited by the presence of Na^{2+} , K^{+} , Mg^{2+} , and Al^{3+} ions. In a previous study, CA activity from *Pseudomonas fragi* was significantly enhanced by Zn^{2+} , Fe^{3+} and Cd^{2+} , Na^{2+} , K^{+} , Mg^{2+} , and Ca^{2+} had no effect, whereas Hg^{2+} and Pb^{2+} had an inhibitory effect [23]. In contrast, the activity of CA from *Dicentrarchus labrax* was inhibited by Zn^{2+} , Hg^{2+} , Al^{3+} , Cu^{2+} , and CO^{2+} [41].

3.5. Enzymatic Conversion of Carbon Dioxide Using Crude Enzyme

Finally, the application of CA to convert CO_2 into CaCO_3 in the presence of calcium ion was examined. Test reaction containing crude CA showed a significant conversion of CO_2 as compared to that of control. BSA was used as a control in the experiment which showed no precipitation of CO_2 . The CO_2 conversion efficacy of CA was studied by calculating the amount of CaCO_3 synthesized. The crude CA resulted in the formation of 45 mg CaCO_3 /mg of protein. The CO_2 conversion efficiency of the enzyme produced from *Corynebacterium flavescens* is much higher as compared to CA from *Bacillus pumilis* (33.06 mg CaCO_3 /mg of protein) and *Pseudomonas fragi* (27.33 mg CaCO_3 /mg of protein) [42, 23].

CaCO_3 is commonly used in various industries such as fillers for paints, papers, plastics, as well as food and pharmaceutical applications [43]. Therefore, it confirmed that the bacterial isolate is producing potent CA which can be used for the CO_2 conversion in the lab as well as commercial scale.

3.6. Instrumental Analysis of CaCO_3

In the present studies, the crude enzyme precipitates CaCO_3 and exhibited morphological resemblance with CaCO_3 standard taken from Sigma Aldrich. In the SEM analysis, spherical vaterite crystal was formed in control and rhombohedral vaterite and calcite crystals were formed in the presence of an enzyme (Fig. 7 (a) and (b)). This result from SEM analysis conforms to the finding of Jo et al. [44]. The composition of precipitated powder was confirmed by XRD analysis. The two forms of CaCO_3 were vaterite, and calcite, analyzed by XRD patterns. Diffraction peak at 29.38, 34.04, 43.90 corresponds to the calcite crystal phase whereas the peak at 22.54, 27.08 corresponds to the vaterite crystal phase (Fig. 7 (c)). Furthermore, various studies have shown that calcite and vaterite were phase formed in the presence of CA [37, 44].

4. Conclusions

Microorganisms are commonly used to produce the enzymes of industrial utility. The isolation of CA from cow saliva is cost-effective as the price of commercially BCA is about \$3,000 g^{-1} . Thus, there

is a price limitation for using BCA in various CO₂ conversion applications. The improved CA production from *Corynebacterium flavesces* was achieved by the optimization of various production and reaction parameters. The high yield of CA was achieved from this process as an indication of the possibility for its large-scale production. The crude enzyme was also found to be effective in the CO₂ conversion experiment and hence its efficacy in CO₂ capture can be exploited at the industrial level too. In future studies, the CA enzyme can be purified and used in an immobilized form for better reaction efficiency. Moreover, directed evolution and rational hybrid design can be used to develop the mutant of this enzyme with better enzyme stability for industrial purposes.

Acknowledgment

The financial support from the Jaypee University of Information Technology, Wanknaghat to undertake this study is thankfully acknowledged. Further, the authors have no conflict of interest either among themselves or with the parent institution.

Author Contributions

T.S. (Ph.D student) performed all the experiments and wrote the manuscript.

A.K. (Assistant Professor) designed experiments and revised the manuscript.

References

- Sharma T, Sharma S, Kamyab H, et al. Energizing the CO₂ utilization by chemo-enzymatic approaches and potentiality of carbonic anhydrases: A review. *J. Clean. Prod.* 2020;247:1-12.
- Zhang W, Liu H, Sun C, et al. Capturing CO₂ from ambient air using a polyethyleneimine-silica adsorbent in fluidized beds. *Chem. Eng. Sci.* 2014;116:306-316.
- Mulenga D, Siziya S. Indoor Air Pollution Related Respiratory Ill Health, a Sequel of Biomass Use. *SciMed. J.* 2019;30-37.
- DiMario RJ, Machingura MC, Waldrop GL et al. The many types of carbonic anhydrases in photosynthetic organisms. *Plant Sci.* 2018;268:11-17.
- Kupriyanova EV, Sinetova MA, Cho SM, et al. CO₂-concentrating mechanism in cyanobacterial photosynthesis: organization, physiological role, and evolutionary origin. *Photosynth Res.* 2013;117(1-3):133-46.
- Chen F, Jin W, Gao H, et al. Cloning, Expression and Characterization of Two Beta Carbonic Anhydrases from a Newly Isolated CO₂ Fixer, *Serratia marcescens* Wy064. *Indian J. Microbiol.* 2018;59(1):64-72.
- Rinanti A. Biotechnology carbon capture and storage by microalgae to enhance CO₂ removal efficiency in closed-system photobioreactor. In: Thajuddin N and Dhanasekaran D eds, *Algae-Organisms for Imminent Biotechnology*. IntechOpen; 2016. p.134-136.
- Hou J, Li X, Kaczmarek MB, et al. Accelerated CO₂ Hydration with Thermostable *Sulfurihydrogenibium azorens* Carbonic Anhydrase-Chitin Binding Domain Fusion Protein Immobilised on Chitin Support. *Int. J. Mol. Sci.* 2019;20(6):1494.
- Chang R, Kim S, Lee S, et al. Calcium carbonate precipitation for CO₂ storage and utilization: A review of the carbonate crystallization and polymorphism. *Front. Energy Res.* 5 2017;1-17.
- Faridi S, Satyanarayana T. Applicability of carbonic anhydrase in mitigating global warming and development of useful products from CO₂. *Clim. Change Environ. Sustain.* 2015;3:77-92.
- Bose H, Satyanarayana T. Utility of thermo-alkali-stable gamma-CA from polyextremophilic bacterium *Aeribacillus pallidus* TSHB1 in biomimetic sequestration of CO₂ and as a virtual peroxidase. *Environ. Sci. Pollut. Res. Int.* 2017;24(11):10869-10884.
- Kumar A, Sharma T, Mulla SI, et al. Let's Protect Our Earth: Environmental Challenges and Implications. In: Kumar A, Sharma S eds. *Microbes and Enzymes in Soil Health and Bioremediation*. Singapore: Springer Singapore; 2019. p. 1-10.
- Aggarwal M, Chua TK, Pinard MA, et al. Carbon Dioxide "Trapped" in a β -Carbonic Anhydrase. *Biochemistry* 2015;54(43):6631-6638.
- Abo-Ashour MF, Eldehna WM, Nocentini A, et al. Novel synthesized SLC-0111 thiazole and thiadiazole analogues: Determination of their carbonic anhydrase inhibitory activity and molecular modeling studies. *Bioorg. Chem.* 2019;87:794-802.
- Alissa SA, Alghulikah HA, Alothman ZA, et al. Phosphoramidates are the first phosphorus-based zinc binding motif to show inhibition of β -class carbonic anhydrases from bacteria, fungi, and protozoa. *J. Enzyme. Inhib. Med. Chem.* 2020;35(1):59-64.
- Boone CD, Habibzadegan A, Gill S, et al. Carbonic anhydrases and their biotechnological applications. *Biomolecules* 2013;3(3):553-62.
- Prete SD, Nocentini A, Supuran CT. Bacterial α -carbonic anhydrase: a new active class of carbonic anhydrase identified in the genome of the Gram-negative bacterium *Burkholderia territorii*. *J. Enzym. Inhib. Med. Ch.* 2020;35(1):1060-1068.
- Hu JJ, Wang L, Zhang SP, et al. Optimization of electron donors to improve CO₂ fixation efficiency by a non-photosynthetic microbial community under aerobic condition using statistical experimental design. *Bioresour. Technol.* 2010;101(18):7073-7078.
- Smith KS, Ferry JG. Prokaryotic carbonic anhydrases. *FEMS Microbiol. Rev.* 2000;24:335-366.
- Anjana G, Dudhagara P, Bhagat C, et al. Biomimetic Sequestration of CO₂ Using Carbonic Anhydrase from Calcite Encrust Forming Marine Actinomycetes. *Sci. Int.* 2015;3:48-57.
- Ramanan R, Kannan K, Sivanesan SD, et al. Bio-sequestration of carbon dioxide using carbonic anhydrase enzyme purified from *Citrobacter freundii*. *World. J. Microb. Biot.* 2009;25(6):981-987.
- Smith KS, Ferry JG. A plant-type (beta-class) carbonic anhydrase in the thermophilic methanarchaeon *Methanobacterium thermoautotrophicum*. *J. Bacteriol.* 1999;181(20):6247-6253.
- Sharma A, Bhattacharya A, Singh S. Purification and characterization of an extracellular carbonic anhydrase from *Pseudomonas fragi*. *Process. Biochem.* 2009;44(11):1293-1297.

24. Sharma A, Sharma T, Meena KR, et al. High throughput synthesis of ethyl pyruvate by employing superparamagnetic iron nanoparticles-bound esterase. *Process. Biochem.* 2018;71:109-117.
25. Meena K, Sharma A, Kumar R, et al. Two factor at a time approach by response surface methodology to aggrandize the *Bacillus subtilis* KLP2015 surfactin lipopeptide to use as anti-fungal agent. *J. King Saud Univ. Sci.* 2018;32(1):337-348.
26. Sharma A, Meena K, Kanwar S. Molecular characterization and bioinformatics studies of a lipase from *Bacillus thermoamylovorans* BHK67. *Int. J. Biol. Macromol.* 2017;107:2131-2140.
27. Giri A, Banerjee UC, Kumar M, et al. Intracellular carbonic anhydrase from *Citrobacter freundii* and its role in bio-sequestration. *Bioresour. Technol.* 2018;267:789-792.
28. Yoshimura H, Iwasaki H, Nishikawa T, et al. Role of carbonic anhydrase in the bicarbonate excretion from salivary glands and mechanism of ionic excretion. *Jpn. J. Physiol.* 1959;9(1): 106-23.
29. Jaya P, Nathan V, Parvathi A. Characterization of marine bacterial carbonic anhydrase and their CO₂ sequestration abilities based on a soil microcosm. *Prep. Biochem. Biotech.* 2019;49:891-899.
30. Kupriyanova E, Villarejo A, Markelova A, et al. Extracellular carbonic anhydrases of stromatolite forming cyanobacterium *Microcoleus cathonoplastes*. *Microbiol.* 2007;153:1149-1156.
31. Buzolyova LS, Somov G. Autotrophic assimilation of CO₂ and C1 compounds by pathogenic bacteria. *Biochemistry* 1996;4 (10):1357-1361.
32. Bose H, Satyanarayana T. Suitability of the alkalistable carbonic anhydrase from a polyextremophilic bacterium *Aeribacillus pallidus* TSHB1 in biomimetic carbon sequestration. *Bioproc. Biosyst. Eng.* 2016;39(10):1515-1525.
33. Shivalee A, Lingappa K, Mahesh D. Influence of bioprocess variables on the production of extracellular chitinase under submerged fermentation by *Streptomyces pratensis* strain KLSL55. *J. Genetic Eng. Biotech.* 2018;16(2):421-426.
34. Zhang Z, Lian B, Hou W, et al. Optimization of nutritional constituents for carbonic anhydrase production by *Bacillus mucilaginosus* K02. *Afr. J. Biotechnol.* 2011;10:8403-8413.
35. Fleuri L, Sato H. Production, purification, cloning and application of lytic enzymes. *Quím. Nova.* 2005;28:871-879.
36. Sundaram S, Thakur IS. Induction of calcite precipitation through heightened production of extracellular carbonic anhydrase by CO₂ sequestering bacteria. *Bioresour. Technol.* 2018;253: 368-371.
37. Li CX, Jiang XC, Qiu YJ, et al. Identification of a new thermostable and alkali-tolerant α -carbonic anhydrase from *Lactobacillus delbrueckii* as a biocatalyst for CO₂ biomineralization. *Bioresour. Bioprocess.* 2015;2(1):44.
38. Chirica LC, Elleby B, Lindskog S. Cloning, expression and some properties of α -carbonic anhydrase from *Helicobacter pylori*. *Biochim. Biophys. Acta.* 2001;1544:55-63.
39. Demir N, Demir Y, Coskun F. Purification and characterization of carbonic anhydrase from human erythrocyte plasma membrane. *Türk. J. Med. Sci.* 2001;31:477-482.
40. Kumar A, Dhar K, Kanwar SS, et al. Lipase catalysis in organic solvents: advantages and applications. *Biol. Proced. Online.* 2016;18:2-2.
41. Ceyhun SB, Senturk M, Yerlikaya E, et al. Purification and characterization of carbonic anhydrase from the teleost fish *Dicentrarchus labrax* (European Seabass) liver and toxicological effects of metals on enzyme activity. *Environ. Toxicol. Pharmacol.* 2011;32:69-74.
42. Prabhu DC, Valechha A, Wanjari S, et al. Carbon composite beads for immobilization of carbonic anhydrase. *J. Mol. Catal. B Enzym.* 2011;71:71-78.
43. Kim IG, Jo BH, Kang DG, et al. Biomineralization-based conversion of carbon dioxide to calcium carbonate using recombinant carbonic anhydrase. *Chemosphere* 2012;87(10):1091-1096.
44. Jo BH, Seo JH, Yang YJ, et al. Bioinspired Silica Nanocomposite with Autoencapsulated Carbonic Anhydrase as a Robust Biocatalyst for CO₂ Sequestration. *ACS Catal.* 2014;4(12): 4332-4340.